TIEMEN WOUTERSEN AND JOHN C. HAM*

UNIVERSITY OF ARIZONA, AND UNIVERSITY OF MARYLAND, IZA, IRP, IFAU, AND IFS

November 2nd, 2015

Abstract. Applied researchers often need to calculate confidence sets for functions of parameters, such as the effects of counterfactual policy changes. If the function is continuously differentiable and has non-zero and bounded derivatives, then they can use the delta method. However, if the function is nondifferentiable (as in the case of simulating functions with zero-one outcomes), has zero derivatives, or unbounded derivatives, then researchers usually use the nonparametric bootstrap or sample from the asymptotic distribution of the estimated parameter vector. Researchers also use these bootstrap approaches when the function is well-behaved but complicated. Indeed, these approaches are advocated by two very influential published articles. We first show that both of these bootstrap methods can produce confidence sets whose asymptotic coverage is less than advertised, i.e. confidence sets that are too small. We then propose two procedures that provide correct coverage asymptotically. In applications, we find that the bootstrap approaches mentioned above produce confidence sets that are significantly smaller than their consistent counterparts, suggesting that previous empirical work is likely to have been overly optimistic in terms of the precision of estimated counterfactual effects.

Keywords: Confidence Sets, simulation, structural models, policy effects. JEL codes: C12, C15.

^{*}Early versions of this paper were titled "Calculating Confidence Intervals for Continuous and Discontinuous Functions of Parameters." Comments are welcome, woutersen@email.arizona.edu and ham@econ.umd.edu. We thank Don Andrews, James Fisher, Bernd Fitzenberger, Ginger Jin, Soohyung Lee, Nicole Lott, Serkan Ozbeklik, Maddie Pickens, Geert Ridder, John Rust, Petra Todd, Yuxin Su, Mauricio Varela, Jonathan Wright and especially Kei Hirano, Xianghong Li, and Chris Sims for helpful comments and discussions. We also thank seminar participants at MIT-Harvard, University of Pennsylvania, University of Wisconsin, UC San Diego, UC Los Angeles, and USC.

1. INTRODUCTION

WE PROPOSE PROCEDURES TO CALCULATE CONFIDENCE SETS for functions of parameters without restricting the derivatives of the functions and without requiring the functions to be continuous. These are the first procedures for these cases that have consistency proofs. The need for such procedures follows from applied work. Applied researchers often calculate confidence sets for functions of estimated parameters, e.g. to carry out counterfactual policy analysis. If the function is differentiable and has non-zero and bounded derivatives, then researchers can use the delta method,¹ although they are often reluctant to use it for complex, nonlinear functions whose derivatives satisfy these properties.

However, if the function has zero or unbounded derivatives, or is discontinuous, as in the case of simulating functions with zero-one outcomes, then the delta method is inappropriate. Krinsky and Robb (1986) propose the following approach as an alternative to the delta method to obtain a $(1 - \alpha)$ confidence set for a function evaluated at the parameter estimates: i) take a large number of draws from the asymptotic (normal) distribution of the parameter estimates; ii) calculate the function value for each draw; and iii) trim ($\alpha/2$) from each tail of the resulting distribution of the function values. Their approach has been widely used in empirical work to obtain confidence sets for complex, nonlinear, differentiable functions of the estimated parameters, such as consumer demand elasticities, the expected duration of unemployment and impulse functions², as well as for nondifferentiable functions of the estimated parameters.³ Finally, two prominent textbooks⁴ also recommend this approach, and the '*wtp*' and '*wtpcikr*' commands in Stata (the leading software package used by applied economists) are based on Krinsky and Robb (1986). Although this procedure of sampling from the asymptotic distribution is sometimes called the parametric bootstrap, this term has more than one meaning in

¹See, e.g., Weisberg (2005) for a description of the delta method.

 $^{^{2}}$ See Krinsky and Robb (1986) and Fitzenberger, Osikominu and Paul (2010) for applications to demand elasticities and unemployment duration respectively. Further, see Inoue and Kilian (2013) for a recent overview of the impulse response function literature. A simple Google search lists forty-four published papers that refer to Krinsky and Robb (1986).

 $^{^{3}}$ A few (of many possible) examples are Gaure, Røed and Westlie (2010), Ham, Mountain and Chan (1997), Hitsch, Hortacsu and Ariely (2010), Merlo and Wolpin (2009) and Røed and Westlie (2011). Its use is advocated, but not implemented, by Eberwein, Ham and LaLonde (2002). A review of the literature indicates that many studies either i) do not give a confidence set for the simulated results or ii) give a confidence set for the simulated results or ii) give a confidence set for the simulated results or ii).

⁴See Greene (2012, page 610) and Wooldridge (2010, page 441).

the literature, so instead we will refer to it as the Asymptotic Distribution bootstrap or AD-bootstrap.

Runkle (1987) recommends the following alternative to the Krinsky-Robb procedure to obtain a $(1-\alpha)$ confidence set for a function evaluated at the parameter estimates: i) draw a bootstrap sample of the data, reestimate the model, and use the resulting parameter estimates to calculate the function; ii) repeat i) many times and trim $(\alpha/2)$ from each tail of the resulting distribution of function values. Runkle's article has been very influential; in fact, it was included in the issue that commemorated the twentieth anniversary of the *Journal of Business and Economics Statistics* as one of the ten most influential papers in the history of the journal (Ghysels and Hall 2002, page 1). Moreover, Runkle's approach is endorsed by three prominent graduate econometrics textbooks.⁵ We refer to this approach as the ADR-bootstrap. It is first order equivalent to the AD-bootstrap for cases where the version of the bootstrap used in Runkle's (1987) procedure estimates the asymptotic distribution of the parameters consistently.

We first give four important examples in which the widely used AD-bootstrap and ADR-bootstrap fail, including one that mimics how researchers construct confidence sets for counterfactual policy analysis via simulation of structural or nonlinear models. We then provide a method of obtaining confidence sets that works in all of these situations under relatively mild conditions (that are likely to be satisfied in empirical work). We also provide a modification of our approach that offers potential efficiency gains in principle and in practice; this second method is asymptotically equivalent to the delta method when the latter is valid. Thus, our proposed procedure is valid under weaker conditions than the delta method but involves no efficiency loss. Therefore, our approach should be very useful in all of the cases where researchers have previously used the AD-bootstrap or the ADR-bootstrap, as well as in the case of differentiable functions where it is unclear whether the (generally numerical) derivatives actually are nonzero and bounded.⁶

⁵Hamilton (1994, page 337), Cameron and Trivedi (2005, page 363) and Wooldridge (2010, page 439); see also Cameron and Trivedi (2010, page 434). Examples of the use of the Runkle (1987) method in applied work are Chaudhuri, Goldberg, and Jia (2006) and Ryan (2012), who use it to obtain confidence intervals for the effects of counterfactual policy changes, and Hoderlein and Mihaleva (2008), who use it to calculate confidence intervals for price elasticities.

 $^{^{6}}$ Hall (1995) provides a bootstrap procedure for calculating standard errors of functions of estimated parameters. However, like the delta method, it does not apply to discontinuous or non-differentiable

To implement our first procedure, the researcher obtains a $(1 - \alpha)$ confidence set for the function of interest by: i) sampling from the asymptotic distribution of the parameter estimator using the bootstrap or using a normal approximation; ii) keeping the draw only if it is in the $(1 - \alpha)$ confidence set for the estimated parameters; iii) calculating the function value for each draw; and iv) using all function values to construct the confidence set for the function. This procedure differs from the AD-bootstrap and ADR-bootstrap in that they trim the extreme values of the function that come from both 'reasonable' values and 'unreasonable' (extreme) values of the parameter vector, while our approach deletes only function values that arise from 'unreasonable' values of the parameters.⁷ We refer to our procedure as the confidence set bootstrap or CS-bootstrap. We also provide a modification that offers potential efficiency gains over the CS-bootstrap, and refer to it as the weighted confidence set bootstrap or the WCS-bootstrap. The substantial conditions that are necessary to apply our approach are: i) that one can sample from the asymptotic distribution of the estimators of the parameters and ii) that the set of points at which the function is discontinuous is small. For example, if the function is a scalar, then the second requirement is that the number of discontinuity points is finite.

We also apply our method to an estimator considered by Andrews (2000). Andrews showed that no version of the bootstrap can consistently estimate the distribution of his maximum likelihood estimator. An example involving a function of a parameter yields the same maximum likelihood estimator as in Andrews (2000). Our proposed procedures also work for this example, suggesting that it might be more fruitful to focus on the construction of confidence sets, rather than on the distributions of various versions of the bootstrap.

A paper related to ours is Hirano and Porter (2012). They show that if the target object is not differentiable in the parameters of the data distribution, then there exists no estimator sequences that is locally asymptotically unbiased or α -quantile unbiased. They note that this places strong limits on estimators, bias correction methods, and inference procedures. Our paper complements this paper in the sense that our method still works

functions bootstrap, but requires stronger assumptions than those necessary for the delta method. Hence we view Hall's procedure as dominated by the delta method and do not consider it further in our paper. ⁷We formalize this notion of 'reasonable' values in Lemma 1.

for a class of models where the function of the parameters is not differentiable in the parameters of the data distribution. Horowitz (2002) states the consistency conditions for the regular bootstrap including the condition that the function of the parameters is differentiable in the parameters of the data distribution. He also gives examples in which this conditions fails.

We use the empirical work from two papers to obtain evidence demonstrating the difference between the procedures in practice. First, we consider work from Ham, Li and Shore-Sheppard (2011, hereafter HLSS), who construct both relatively simple differentiable functions, and relatively complicated nondifferentiable functions, of their parameter estimates describing the labor market dynamics of disadvantaged women in the U.S. Second, we consider confidence sets for complex differentiable functions of estimated parameters from a rich model of dating and marriage that Lee and Ham (2012, hereafter LH) use to evaluate the efficacy of different matching mechanisms for online dating. We find first that for HLSS' simple differentiable functions, the AD-bootstrap produces somewhat smaller confidence sets than the (appropriate) delta method. Second, we find that the AD-bootstrap produces much smaller confidence sets than those from the (appropriate) CS-bootstrap for LH's complicated differentiable functions and HLSS' nondifferentiable functions. Additionally, we find that the WCS-bootstrap offers substantial efficiency gains over the CS-bootstrap in the case of relatively simple differentiable functions. The upshot is that the size of many estimated confidence sets in the literature may be substantially biased downwards.

We proceed as follows. In section 2 we show that, in several important examples, the AD-bootstrap and ADR-bootstrap fail to provide a confidence set with the correct asymptotic coverage. In section 3 we show that the CS-bootstrap and the WCS-bootstrap provide consistent confidence sets for both nondifferentiable and discontinuous functions. In section 4 we provide evidence on the difference between the CS, WCS and AD-bootstraps in practice, and we conclude in section 5.

2. Failures of the Delta Method and the AD-bootstrap when Calculating Confidence Sets for Functions of Parameters

We examine the performance of the delta method, the AD-bootstrap, and the ADRbootstrap in four important examples in this section. However, since the AD-bootstrap and ADR-bootstrap are equivalent asymptotically, for expositional ease in what follows we simply refer to the AD-bootstrap with the understanding that the ADR-bootstrap will behave in exactly the same way as the AD-bootstrap. In Example 1, both the delta method and the AD-bootstrap fail. In Example 2, the AD-bootstrap fails. In Example 3, the delta method is infeasible and no version of the bootstrap consistently estimates the asymptotic distribution of the function of the estimator; however, we show below that the CS-bootstrap can be used to construct a valid confidence set. Example 4 will be of most interest to applied researchers, since it mimics how researchers construct confidence sets for counterfactual policy analysis where outcomes are discrete and policy effects are obtained via simulation of structural or nonlinear models. However, it is also our most complicated example, and our first three examples, while somewhat more theoretical in nature, should prove useful to even very applied researchers. The outcomes here are discontinuous functions of the data, and researchers have turned to the AD bootstrap because the delta method is not applicable. However, we show that the AD bootstrap again fails in this very important application.

Example 1: The Delta Method and the AD-bootstrap Fail

Suppose we observe a random sample, $X_1, ..., X_N$, from a normal distribution with mean μ and variance 1, and let $\hat{\mu} = \bar{X}_N = \frac{1}{N} \sum_i X_i$. Let $E(X) = \mu_0 = 0$ and consider $h(\mu) = \sqrt{|\mu|}$. The delta method yields the following symmetric 95% confidence interval with probability one:

$$\left[\sqrt{|\bar{X}_N|} - \frac{1.96}{2\sqrt{N}\sqrt{|\bar{X}_N|}}, \sqrt{|\bar{X}_N|} + \frac{1.96}{2\sqrt{N}\sqrt{|\bar{X}_N|}}\right]$$

The probability that the true value is inside this confidence interval is about 0.67 (in repeated samples) for any N. Fortunately, our method gives a confidence interval with the correct coverage probability of 95%.

In Example 1, the delta method fails because the derivative does not exist at one

point. The failure is easy to spot here. However, such problems may be much harder to spot in more realistic applications such as the two empirical applications that we consider below, both of which involve estimating more than one hundred parameters. In example 1, the AD-bootstrap also fails. In particular, it yields a confidence set with a coverage probability of 0% for any N. The reason for this is that the true value, $h(\mu = 0) = 0$, is the minimum of the function. Let G(.) denote the distribution function of $h(\hat{\mu})$. Applying the AD-bootstrap and using the interval between the 2.5 and 97.5 percentiles of the function values, [G(0.025), G(0.975)], yields a confidence set that does *not* cover the true value, so the coverage probability is 0%.

Example 2: A Probit Model

Suppose that one is interested in the function $h(\beta, \gamma) = \frac{1}{2}\Phi(\beta) + \frac{1}{2}\Phi\left(-2\gamma - \sqrt{2\ln(2)}\right)$ where the true values of the parameters are zero, i.e. $\beta_0 = \gamma_0 = 0$. Let the estimator $(\hat{\beta}, \hat{\gamma})$ have a normal distribution with mean zero and a known variance-covariance matrix $\begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix}$. The delta method cannot be used since the function has a zero derivative at the true value of the parameters if $\rho = 1$. The AD-bootstrap samples from the normal distribution with mean $(\hat{\beta}, \hat{\gamma})$ and covariance matrix $\begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix}$ in order to construct the distribution of $h(\hat{\beta}, \hat{\gamma})$. Let G(.) denote this distribution function of $h(\hat{\beta}, \hat{\gamma})$. Applying the AD-bootstrap and using the interval between the 2.5 and 97.5 percentiles of the function values, [G(0.025), G(0.975)], does not yield a confidence set with 95% coverage for many values of ρ . For example, the coverage is 0% for the AD-bootstrap if $\rho = 1,90\%$ for $\rho = 0.5$, and 93% for $\rho = 0.8$ Thus, the AD-bootstrap does not produce a confidence set with the correct coverage.⁹ We also note that the AD-bootstrap confidence set coincides with the Bayesian credible interval (with flat priors) in this case, so the Bayesian procedure also fails here. Note that the extreme failure of the AD-bootstrap for $\rho = 1$ occurs because $h(\beta_0 = 0, \gamma_0 = 0) = 0$ is the minimum value of the function.¹⁰ Just as in example 1, this is the case even if the variance of the estimators is arbitrarily small. We observe that the continuity of the coverage as a function of the true values of the parameters also causes

⁸See the Appendix for details.

⁹Correct coverage of a confidence set means that the coverage probability is no smaller than its nominal probability, see Andrews and Cheng (2012 and 2013).

¹⁰Thus, sampling from the asymptotic distribution yields values of the function that are larger than the minimum (and true value) with probability one, ensuring failure of the AD-bootstrap.

the AD-bootstrap to fail for ρ close to one. Finally, we note that using the AD-bootstrap to calculate standard errors for average partial effects can also fail in this type of situation.

Example 3: Andrews (2000), Inconsistency of the Bootstrap

Suppose we observe a random sample, $X_1, ..., X_N$, from a normal distribution with mean μ and variance 1 (denoted as $N(\mu, 1)$) and suppose that μ is restricted to be nonnegative. Andrews (2000) considers the maximum likelihood estimator $\hat{\mu} = \max(\bar{X}_N, 0)$ where $\bar{X}_N = \frac{1}{N} \sum_i X_i$. He shows that the nonparametric bootstrap fails to consistently estimate the distribution of $\hat{\mu}$ if $\mu = 0$, and that it is impossible to consistently estimate the distribution of $\hat{\mu}$ using any bootstrap method if $\mu = \frac{c}{\sqrt{N}}$ for any c > 0. A related problem is deriving a confidence set for the function $h(\gamma) = \max(\gamma, 0)$ where $\gamma = E(X)$ and we observe a random sample, $X_1, ..., X_N$, from a normal distribution with mean γ and variance 1. Note that γ can be negative and let $\hat{\gamma} = \bar{X}_N$. The estimator $h(\hat{\gamma}) = \max(\bar{X}_N, 0)$ is the identical to Andrews' (2000) estimator and it follows from the reasoning of Andrews (2000) that no version of the bootstrap can estimate the distribution of $h(\hat{\gamma})$ if $\gamma = \frac{c}{\sqrt{N}}$ for any c > 0. We show that in this case the CS-bootstrap can consistently estimate the confidence set for $h(\gamma)$.

The last example considers a case that uses simulations to estimate the function of interest.

Example 4: An Ordered Probit Model

Suppose that a firm offers three products, A, B, and C, where product B is more luxurious than product A, but less luxurious than product C. Let consumer i make the following choice:

> $A_i = 1$ if $-2 \le X_i\beta + \varepsilon_i < 0$ and $A_i = 0$ otherwise, $B_i = 1$ if $0 \le X_i\beta + \varepsilon_i \le 2$ and $B_i = 0$ otherwise, and $C_i = 1$ if $X_i\beta + \varepsilon_i > 2$ and $C_i = 0$ otherwise,

where X_i equals one if the consumer lives in a market where the firm is advertising and $\varepsilon_i | X_i$ has a standard normal distribution. Note that consumer *i* does not make a purchase if $X_i\beta + \varepsilon_i < -2$. Suppose that the firm conducted an advertising campaign in one market to encourage customers to buy a more luxurious product and that the firm estimated β . Next, an econometrician's objective is to predict how the demand for product *B* responds to such advertising in another market, including a confidence set for this response. Suppose that the true value of β equals one and that the estimator for β is normally distributed with mean 1 and variance $\sigma_{\beta}^2 = 1$. In such a case, it is natural to estimate the change in the demand for product *B* using simulations. This is a case where the AD-bootstrap fails. In particular, we used simulated data on 1000 individuals and simulated 100 draws from the distribution of ε for each individual.¹¹ We then drew 1,000 times from the distribution of the estimator and calculated the change in the probability of buying good *B*. The resulting AD-bootstrap 95% confidence set covered only the true value of this change, 0.2054, in 36% of the cases.¹² Simulating data to form confidence sets often yields non-differentiable functions, as in this example. Therefore, the delta method cannot be used, while the AD-bootstrap may yield inconsistent results. Our applied section presents more complicated examples that use simulations.

Note that subsampling, as proposed by Politis and Romano (1994), does not yield confidence sets with asymptotically correct coverage in the examples above. Of course, there are examples where the AD-bootstrap will have correct asymptotic coverage, but it is difficult to ascertain in general when this will be the case.¹³ In particular, Hirano and Porter (2012) show that if the target object is not differentiable in the parameters of the data distribution, then there exist no estimator sequences that are locally asymptotically unbiased or α -quantile unbiased. They then note "Since no regular estimator exists, the usual arguments for the validity of standard approaches to inference, such as Wald-type procedures, will not be valid." The examples above show that it can be difficult to see

¹¹Some readers may question the need to use the AD bootstrap, since with a very large number of simulations the outcome may approach a continuously differentiable function and researchers should be able to use the delta method. However, our example is a toy model, and using a very large number of simulations in a function evaluation is not feasible for the structural models used for counterfactual policy analysis.

¹²Also, the coverage probability remained well below 95%, even for much smaller values of σ_{β}^2 .

¹³For example, it is straightforward to show that if θ and $h(\theta)$ are scalars, then a sufficient condition for the AD-bootstrap to work is for $h(\theta)$ to be monotonic. However, example 2 shows that this does not generalize to the case where the parameter is of dimension two and the function is monotonic in its first and second argument.

whether the "target object" is differentiable in the parameters of the data distribution so that it is desirable to have a procedure that yields valid confidence sets even if this differentiability is violated.

3. MAIN RESULT

In this section we provide a method of obtaining confidence sets that is valid under reasonable assumptions that are likely to be satisfied in empirical work. We begin by discussing the CS-bootstrap and then consider the WCS-bootstrap. The latter has the advantage that it is asymptotically equivalent to the delta method when the delta method is valid. For both of these approaches, we consider using the asymptotic distribution and the bootstrap approximation.

Let the dimension of θ be equal to K and let $h(\theta)$ have dimension H. Note that allowing for H > 1 is important, since it allows one to obtain a joint confidence set for multiple counterfactual outcomes. For example, in a structural model with human capital accumulation (Eckstein and Wolpin 1999, Keane and Wolpin 2000), one can look at the effect of a policy change on both completed schooling and work experience (at any point in the life cycle). Alternatively, in a model of labor market dynamics (Eberwein, Ham and LaLonde 1997), it would be helpful to have a joint confidence set for the effect of participating in a training program on the expected duration of employment and the expected duration of unemployment.

Suppose that the estimator for θ , denoted by $\hat{\theta}$, is asymptotically normally distributed and consider the following confidence set for the parameter θ :

$$CS^{\theta}_{1-\alpha} = \{\theta \in \Theta | N \cdot (\hat{\theta} - \theta)'(\hat{\Omega})^{-1}(\hat{\theta} - \theta) \le \chi^2_{1-\alpha}(K)\},\tag{1}$$

where $\hat{\Omega}$ is the asymptotic variance-covariance matrix for $\hat{\theta}$ and $\chi^2_{1-\alpha}(K)$ is the $(1-\alpha)$ percentile of the χ^2 distribution with K degrees of freedom. Next, let $CS_{1-\alpha}^{h(\theta)}$ denote the set of values that we obtain if we apply the function $h(\theta)$ to every element of $CS_{1-\alpha}^{\theta}$. More precisely,

$$CS_{1-\alpha}^{h(\theta)} = \{ \tau \in \mathbb{R}^H | \tau = h(\theta) \text{ for some } \theta \in CS_{1-\alpha}^{\theta} \}.$$
 (2)

Suppose that the researcher draws M^* times from the asymptotic distribution of $\hat{\theta}$. Let $\tilde{\theta}_1, ..., \tilde{\theta}_{M^*}$ denote these draws. The researcher then keeps the draws that satisfy $\tilde{\theta}_m \in CS^{\theta}_{1-\alpha}, m = 1, ..., M^*$.. Let $\tilde{\theta}_1, ..., \tilde{\theta}_M$ denote retained these draws. We now estimate the confidence set for the function of the parameters, $CS^{h(\theta)}_{1-\alpha}$, by applying the function h(.) to the draws $\tilde{\theta}_1, ..., \tilde{\theta}_M$. In particular, let $CS^{h(\theta)}_{1-\alpha}$ be the set of all points in the image of $h(\theta), \theta \in \Theta$, that are no farther than the Euclidian distance $\eta > 0$ away from $h(\tilde{\theta}_1), ..., h(\tilde{\theta}_{M-1})$, or $h(\tilde{\theta}_M)$.¹⁴ The confidence set $CS^{h(\theta)}_{1-\alpha}$ is what arises from using what we described heuristically above as the CS-bootstrap.

We briefly note why the AD-bootstrap can fail for the case where $h(\theta)$ is a scalar. The AD-bootstrap samples from the entire asymptotic distribution of $\hat{\theta}$ and forms the confidence set of $h(\theta)$ by trimming the extreme $(1-\alpha)/2$ values from the upper and lower tails of the resulting distribution for $h(\theta)$. Note that the extreme values of $h(\theta)$ that the AD-bootstrap trims can arise from i) an extreme draw from the asymptotic distribution of θ or ii) a 'reasonable' draw for θ that results in an extreme value of $h(\theta)$.¹⁵ The CS-bootstrap instead samples from the $(1 - \alpha)$ confidence set of θ and includes all of the resulting values of $h(\theta)$ in its $(1 - \alpha)$ confidence set, and thus does not trim $h(\theta)$ for a 'reasonable' draw of θ . Moreover, note that constructing a confidence set using the CS-bootstrap is no more difficult than constructing one using the AD-bootstrap.

A similar procedure can be used if the researcher draws J bootstrap samples to obtain the distribution of $\hat{\theta}$. Let $\tilde{\theta}_1, ..., \tilde{\theta}_J$ denote the bootstrap sample estimates and $\hat{\Omega}$ the variance-covariance matrix of the bootstrap samples. For each estimate, we calculate $B_j = (\hat{\theta} - \tilde{\theta}_j)'\hat{\Omega}^{-1}(\hat{\theta} - \tilde{\theta}_j), j = 1, ..., J$. We then select the $(1 - \alpha) \cdot J$ bootstrap estimates that have the smallest values of B_j , and call this set B. Let $\tilde{\theta}_1, ..., \tilde{\theta}_M$ denote these draws. As before, we estimate the confidence set by applying the function h(.) to the draws $\tilde{\theta}_1, ..., \tilde{\theta}_M$. That is, $\widehat{CS_{1-\alpha}^{h(\hat{\theta})}}$ is the set of all points in the image of $h(\theta), \theta \in \Theta$, that are no farther than the Euclidian distance $\eta > 0$ away from $h(\tilde{\theta}_1), ..., h(\tilde{\theta}_{M-1})$, or $h(\tilde{\theta}_M)$.

¹⁴Let Θ_h be the image of $h(\theta)$, $\theta \in \Theta$. If $\tilde{\theta}_s \in CS^{\theta}_{1-\alpha}$ is sampled, then any $h \in \Theta_h$ for which $||h(\tilde{\theta}_s) - h||^2 \leq \eta$ is included in the $(1 - \alpha)$ confidence set for $h(\theta)$. If $CS^{h(\theta)}_{1-\alpha}$ is convex then it may be possible to calculate the set directly without using η , e.g. $h(\theta)$ is a scalar and $\{\underline{h}=\min h(\theta): \theta \in CS_{1-\alpha}\}$ and $\{\overline{h}=\max h(\theta): \theta \in CS_{1-\alpha}\}$ can be calculated so that $CS^{h(\theta)}_{1-\alpha} = [\underline{h}, \overline{h}]$.

¹⁵In the lemma that follows, we formalize the notion that values that are closer to $\hat{\theta}$ are likely to be closer to the true value θ_0 as well (compared to values that are further away from $\hat{\theta}$).

We next consider the case where the asymptotic distribution of $\hat{\theta}$ is unknown but one can construct a confidence set for it. For example, $CS_{1-\alpha}^{\theta}$ is derived using bounds.¹⁶ Note that one cannot calculate $h(\theta)$ for every $\theta \in CS_{1-\alpha}^{\theta}$. Therefore, we use a grid that has Mpoints to approximate $CS_{1-\alpha}^{\theta}$. We then calculate $h(\theta)$ for each of these M grid points.¹⁷ Next, let the confidence set $\widehat{CS_{1-\alpha}^{h(\theta)}}$ be the set of all points in the image of $h(\theta), \theta \in \Theta$, that are no farther than the Euclidian distance $\eta > 0$ away from $h(\tilde{\theta}_1), ..., h(\tilde{\theta}_{M-1})$, or $h(\tilde{\theta}_M)$.

We state our first assumption in terms of properties of $CS_{1-\alpha}^{\theta}$. Later on, when we discuss the weighted CS-bootstrap, we use properties of the asymptotic distribution as primitives in the assumptions, since the weighting may depend on this asymptotic distribution. Let N denote the sample size. Also, let P be the data generating process and let \mathcal{P} be a space of probability distributions. Our first assumption requires the true value of the parameter, $\theta_0(P)$, to be an element of $CS_{1-\alpha}^{\theta}$ with probability of at least $(1 - \alpha)$, uniformly over \mathcal{P} .

Assumption 1

Let (i) $\theta_0(P) \in \Theta$, which is compact; and (ii)

$$\lim_{N \to \infty} \inf_{P \in \mathcal{P}} \Pr\{\theta_0(P) \in CS^{\theta}_{1-\alpha}\} \ge 1 - \alpha$$

for any $\alpha \in (0, 1)$.

While the true parameter, $\theta_0(P)$, is of course a function of the data generating process, for expositional ease we often write it as θ_0 . Note that Assumption 1 simply says that the confidence set for the parameter contains the true parameter value with probability $(1 - \alpha)$ in the limit, uniformly over \mathcal{P} . This will certainly hold for any estimator that is uniformly asymptotically normally distributed, as well as for the subsampling and bootstrap confidence sets for θ under appropriate regularity conditions (see Romano and Shaikh 2010, and Andrews and Guggenberger 2010).

 $^{^{16}}$ E.g. the confidence set is derived using the techniques proposed by Imbens and Manski (2004), Chernozhukov, Hong, and Tamer (2007), or Stoye (2009).

¹⁷One can use equally spaced grids, Halton sequences, Halton (1964), or Sobol sequences, Sobol (1967). All these grids are dense in $CS_{1-\alpha}^{\theta}$ as M increases.

Assumption 2

Let $h(\theta)$ be bounded over the domain Θ . Let there exist a partitioning of the parameter space such that $\Theta_1 \cup \Theta_2 ... \cup \Theta_R = \Theta$, where $R < \infty$; let $\Theta_1, \Theta_2, ..., \Theta_{R-1}$ and Θ_R have nonzero Lebesgue measure; and let $h(\theta)$ be uniformly continuous¹⁸ for all $\theta \in \Theta_r$, r = 1, ..., R.

The second assumption allows $h(\theta)$ to be discontinuous. For example, if θ is a scalar, then Assumption 2 requires that the number of discontinuities is finite. In general, the parameter space is partitioned into R subsets, and $h(\theta)$ is assumed to be uniformly continuous on each of these sets. The restriction is that R is finite. This condition is weaker than the conditions needed for the delta method.

Next, we propose a modified version of our procedure. This modified procedure uses weights and usually yields a smaller confidence set than the CS-bootstrap. The idea is to use a weighted average of the elements of the parameter vector θ . These weights are comparable to the weights used in the general method of moments (GMM) procedure, in the sense that the reason to use them is to reduce variation or spread. For example, consider the function $h(\theta) = \Phi(\theta_1 + 2\theta_2)$; then the researcher could use a confidence set for $\theta_1 + 2\theta_2$ rather than the confidence set for (θ_1, θ_2) . That is, the researcher could use a confidence set for a weighted average. In general, let $\hat{\theta}$ be asymptotically normally distributed and let $\hat{\Omega}$ denote a consistent estimator for its asymptotic variance-covariance matrix. Define the vector $w = (w_1, w_2, ..., w_K)'$, where $w_1, w_2, ..., w_K$ are scalars if $h(\theta)$ is a scalar and column vectors with length H otherwise. Consider the following confidence set for $w'\theta$:

$$WCS^{\theta}_{1-\alpha} = \{\theta \in \Theta | N \cdot (\hat{\theta} - \theta)' w (w' \hat{\Omega} w)^{-1} w' (\hat{\theta} - \theta) \le \chi^2_{1-\gamma}(H)$$

and $N \cdot (\hat{\theta} - \theta)' \hat{\Omega}^{-1} (\hat{\theta} - \theta) \le \chi^2_{1-\frac{\gamma}{5}}(K) \}.$

If $h(\theta)$ is a scalar, as in the applications reviewed in the Introduction, then H = 1. Let $WCS_{1-\alpha}^{h(\theta)}$ denote the set of values that we obtain if we apply the function $h(\theta)$ to every

¹⁸The vector-function $h(\theta)$ is uniformly continuous on Θ_j if for any $\eta > 0$ there is an $\varepsilon > 0$ such that $||h(\theta_1) - h(\theta_2)|| < \eta$ for all $\theta_1, \theta_2 \in \Theta_j$ with $||\theta_1 - \theta_2|| < \varepsilon$ where ||.|| is the Euclidean norm.

element of $WCS^{\theta}_{1-\alpha}$. That is,

$$WCS_{1-\alpha}^{h(\theta)} = \{ \tau \in \mathbb{R}^H | \tau = h(\theta) \text{ for some } \theta \in WCS_{1-\alpha}^{\theta} \}.$$
 (3)

We estimate $WCS_{1-\alpha}^{h(\theta)}$ by drawing M^* times from the asymptotic distribution of $\hat{\theta}$, and keeping the draws that are elements of $WCS_{1-\alpha}^{\theta}$ (i.e. draws that satisfy $N \cdot (\hat{\theta} - \theta)'w(w'\hat{\Omega}w)^{-1}w'(\hat{\theta}-\theta) \leq \chi_{1-\gamma}^2(H)$ and $N \cdot (\hat{\theta}-\theta)'\hat{\Omega}^{-1}(\hat{\theta}-\theta) \leq \chi_{1-\frac{\gamma}{5}}^2(K)$). Let $\tilde{\theta}_1, ..., \tilde{\theta}_M$ denote these retained draws. We now estimate this confidence set by applying the function h(.) to the draws $\tilde{\theta}_1, ..., \tilde{\theta}_M$. In particular, let $\widehat{WCS_{1-\alpha}^{h(\theta)}}$ be the set of all points in the image of $h(\theta), \theta \in \Theta$, that are no farther than the Euclidian distance $\eta > 0$ away from $h(\tilde{\theta}_1), ..., h(\tilde{\theta}_{M-1})$, or $h(\tilde{\theta}_M)$.

A similar procedure can be used if the researcher uses the bootstrap for the distribution of $\hat{\theta}$. Again, let $\tilde{\theta}_1, ..., \tilde{\theta}_J$ denote the bootstrap sample estimates and $\hat{\Omega}$ the variancecovariance matrix of the bootstrap samples. For each estimate, we calculate $\tilde{B}_j = (\hat{\theta} - \tilde{\theta}_j)'w(w'\hat{\Omega}w)^{-1}w'(\hat{\theta} - \tilde{\theta}_j), j = 1, ..., J$. We then select the $(1 - \gamma) \cdot J$ bootstrap estimates that have the smallest values of \tilde{B}_j and call this set \tilde{B} . Next we select the $(1 - \alpha) \cdot J$ bootstrap estimates that have the smallest values of $\bar{B}_j = (\hat{\theta} - \tilde{\theta}_j)'\hat{\Omega}^{-1}(\hat{\theta} - \tilde{\theta}_j), j = 1, ..., J$. and call this set \bar{B} .Let $\tilde{\theta}_1, ..., \tilde{\theta}_M$ denote these draws. As before, we estimate the confidence set by applying the function h(.) to the draws $\tilde{\theta}_1, ..., \tilde{\theta}_M$. That is, $WCS_{1-\alpha}^{h(\hat{\theta})}$ is the set of all points in the image of $h(\theta), \theta \in \Theta$, that are no farther than the Euclidian distance $\eta > 0$ away from $h(\tilde{\theta}_1), ..., h(\tilde{\theta}_{M-1})$, or $h(\tilde{\theta}_M)$.

In applications, the weights w will often be estimated. One may estimate w by using numerical derivatives of $h(\theta)$ around the estimate $\hat{\theta}$. The numerical derivatives provide simple estimates for the weights, \hat{w} , and then one replaces w by \hat{w} in forming $\widehat{WCS_{1-\alpha}^{\theta}}$ and \tilde{B} to obtain confidence sets for $h(\theta)$. Furthermore, we suggest limiting the ratio of the weights so that $\min_k(|\hat{w}_k|)/\max_k(|\hat{w}_k|) \geq 1/100$. The WCS-bootstrap yields confidence sets with the correct asymptotic coverage for $h(\theta)$, even if some of the partial derivatives of $h(\theta)$ are infinite (as in Example 1) or zero, while of course this is not true for the delta method. Since the WCS-bootstrap is asymptotically equivalent to the delta method when the latter is valid (see the Appendix), the WCS-bootstrap is safer to use than the delta method but involves no loss of efficiency.

A somewhat more complicated procedure that avoids numerical differentiation is the following. First, consider the case where the researcher samples from the asymptotic distribution. In that case, we propose letting the initial confidence set be all values of $\theta \in \Theta$ for which $N \cdot (\hat{\theta} - \theta)' \hat{\Omega}^{-1} (\hat{\theta} - \theta) \leq \chi^2_{1-\alpha}(K)$. We then use the values of this initial confidence set to estimate a linear approximation to the function $h(\theta)$. In particular, we use the asymptotic distribution to draw M values of the parameter that satisfy $N \cdot (\hat{\theta} - \theta)' \hat{\Omega}^{-1} (\hat{\theta} - \theta) \leq \chi^2_{1-\alpha}(K)$. Let $\tilde{\theta}_1, ..., \tilde{\theta}_M$ denote these points. We then calculate $h(\tilde{\theta}_1), ..., h(\tilde{\theta}_M)$ and regress $\{h(\tilde{\theta}_1), ..., h(\tilde{\theta}_M)\}$ on $\{\tilde{\theta}_1, ..., \tilde{\theta}_M\}$. Let \hat{w} denote the least

late $h(\tilde{\theta}_1), ..., h(\tilde{\theta}_M)$ and regress $\{h(\tilde{\theta}_1), ..., h(\tilde{\theta}_M)\}$ on $\{\tilde{\theta}_1, ..., \tilde{\theta}_M\}$. Let \hat{w} denote the least squares estimator and use the elements of \hat{w} as weights. Note that $\hat{w}'\theta$ is just the best linear predictor and that again $h(\theta)$ is not required to be continuous. Next, construct a confidence set for $h(\theta)$ by again replacing w with \hat{w} .

A similar procedure can be used to estimate weights if the researcher uses the bootstrap. Once again, let $\tilde{\theta}_1, ..., \tilde{\theta}_J$ denote the bootstrap sample estimates and $\hat{\Omega}$ the variancecovariance matrix of the bootstrap samples. As in the case of the CS-bootstrap, we calculate $\tilde{A}_j = (\hat{\theta} - \tilde{\theta}_j)'\hat{\Omega}^{-1}(\hat{\theta} - \tilde{\theta}_j), j = 1, ..., J$. We then select the $(1 - \alpha) \cdot J$ bootstrap estimates that have the smallest values of \tilde{A}_j and call this set \tilde{A} . Next, we regress $h(\tilde{\theta}_j)$ on $\tilde{\theta}_j$ using all $j \in \tilde{A}$. This yields the weights \hat{w} . Next, construct a confidence set for $h(\theta)$ by again replacing w with \hat{w} .

Besides Assumption 2, we also need Assumption 3 for the WCS-bootstrap when we construct $\widehat{WCS}_{1-\alpha}^{h(\theta)}$ (i.e. sample from the asymptotic distribution of $\hat{\theta}$).

Assumption 3

Let (i) $\theta \in \Theta$, which is compact; (ii) for all $k, w_k \neq 0, \hat{w}_k \neq 0, \sup_{P \in \mathcal{P}} |\hat{w}_k - w_k| = o_p(1);$ (iii) $\sqrt{N}(\hat{\theta} - \theta_0) \xrightarrow{d} N(0, \Omega)$ uniformly in $P \in \mathcal{P}$, and the estimator $\hat{\Omega}$ converges to Ω uniformly in $P \in \mathcal{P}$, where Ω and $w'\Omega w$ have full rank; (iv) $\alpha \in (0, 1)$.

If the researcher uses the bootstrap to obtain the confidence set for θ , then we need an additional assumption for $\widehat{WCS_{1-\alpha}^{h(\theta)}}$. In particular, we require that the weighted average, $\widehat{w}'\theta_0$, is in the confidence set $WCS_{1-\alpha}^{\theta}$ with a probability that is equal to or larger than $(1-\alpha)$, uniformly in $P \in \mathcal{P}$. Romano and Shaikh (2010) give uniform convergence results for the bootstrap (and subsampling).

Assumption 4

If a version of the bootstrap is used, then

$$\lim_{N \to \infty} \inf_{P \in \mathcal{P}} \Pr(\hat{w}' \theta_0 \in WCS^{\theta}_{1-\alpha}) \ge 1 - \alpha.$$

Before stating our theorem, intuition for our result can be obtained by continuing our consideration of example 3.

Example 3 (Continuation):

Consider $h(\gamma) = \max(\gamma, 0)$ where $\gamma = E(X)$ and $\hat{\gamma} = \bar{X}_N$. It follows from Andrews that the nonparametric bootstrap fails to consistently estimate the distribution of $\hat{\mu}$ if $\mu = 0$. Further, he demonstrates that it is impossible to consistently estimate (using any version of the bootstrap) the distribution of $\hat{\gamma}$ if $\gamma = \frac{c}{\sqrt{N}}$ for any c > 0. However, the CS-bootstrap can be used to calculate a 95% confidence set (or a $(1 - \alpha)$ confidence set) for $h(\gamma)$ in spite of the absence of a consistent estimator of the asymptotic distribution. In particular, let $\delta = h(\gamma) = \max(\gamma, 0)$. The symmetric 95% confidence set for δ is $\left[\max\left(0, \bar{X}_N - \frac{1.96}{\sqrt{N}}\right), \max\left(0, \bar{X}_N + \frac{1.96}{\sqrt{N}}\right)\right]$, which contains δ with probability 0.95, including the case where $\delta = \frac{c}{\sqrt{N}}$ for any c > 0.¹⁹

We now state our theorem.

Theorem

Let Assumptions 1-2 hold. Then the CS-bootstrap yields

$$\lim_{M \to \infty} \lim_{N \to \infty} \inf_{P \in \mathcal{P}} \Pr\left(h(\theta_0) \in \widehat{CS_{1-\alpha}^{h(\theta)}}\right) \ge 1 - \alpha.$$

Let Assumptions 2-3 hold. Then sampling from the asymptotic distribution and using the WCS-bootstrap yields

$$\lim_{M \to \infty} \lim_{N \to \infty} \inf_{P \in \mathcal{P}} \Pr\left(h(\theta_0) \in \widehat{WCS_{1-\alpha}^{h(\theta)}}\right) \ge 1 - \alpha.$$

Let Assumptions 2-4 hold. Then using a bootstrap procedure for $\hat{\theta}$ and the WCS-bootstrap yields

$$\lim_{M \to \infty} \lim_{N \to \infty} \inf_{P \in \mathcal{P}} \Pr\left(h(\theta_0) \in \widehat{WCS_{1-\alpha}^{h(\theta)}}\right) \ge 1 - \alpha.$$

 $^{^{19}}$ This example also illustrates that one should perhaps not focus exclusively on the distribution of the bootstrap when the goal is to derive a confidence set. Also, Hirano and Porter (2012) derive more impossibility results.

Proof: See Appendix.

If one relaxes the uniformity requirements²⁰ in Assumption 1, 3, or 4, then the theorem holds without the uniformity property. Specifically, if we replace Assumption 1 by (i) $\theta_0 \in \Theta$, which is compact; and (ii)

$$\lim_{N \to \infty} \Pr(\theta_0 \in CS^{\theta}_{1-\alpha}) \ge 1 - \alpha,$$

then the theorem holds without the uniformity result, i.e.

$$\lim_{M \to \infty} \lim_{N \to \infty} \Pr\left(h(\theta_0) \in \widehat{CS_{1-\alpha}^{h(\theta)}}\right) \ge 1 - \alpha.$$

Also, Assumption 3 puts only mild restrictions on the weights. In particular, one could use other estimators for the weights. For example, if the function $h(\theta)$ has a single index, then one also could calculate the weights using semiparametric least squares, as in Ichimura (1993), or one of the single-index estimators reviewed by Horowitz (1998). In general, our weighting is analogous to the use of a weighting matrix when applying the method of moments estimator. In particular, using a weighting matrix that does *not* converge to the efficient weighting matrix does not, in general, cause the method of moments estimator to be inconsistent, see Hansen (1982) and Newey and McFadden (1994). The same is true here for the choice of weights, \hat{w}_k , k = 1, ..., K. Choosing an efficient weighting matrix is, in general, a good idea and here we suggest using the WCS-bootstrap with nonzero weights rather than the CS-bootstrap. Using nonzero weights is analogous to the approach of Newey and West (1987) and Andrews (1991), who advocate using estimates of the variance-covariance matrix that are positive semi-definite.

The main difference between the CS and WCS-bootstrap on the one hand and the AD-bootstrap on the other hand is that the CS and WCS-bootstrap use values of θ that are close to $\hat{\theta}$, while the AD-bootstrap does not have this property. In particular, the AD-bootstrap trims extreme values of $h(\theta)$ rather than extreme values of θ . This explains why the AD-bootstrap yields an inconsistent confidence set in Example 2. We formalize the notion that values of θ that are closer to $\hat{\theta}$ also are likely to be closer to the true value θ_0 in the following lemma.

²⁰Andrews (1987) emphasizes the importance of uniform convergence.

Lemma

Let θ , v, and w be scalars. Let $(\hat{\theta} - \theta_0) \sim N(0, \sigma^2), \sigma^2 > 0$, and $v^2 < w^2$. Then

 $P(|\hat{\theta} + v - \theta_0| \le u) > P(|\hat{\theta} + w - \theta_0| \le u) \text{ for any } u > 0.$

Proof: See appendix.

The lemma states that $\hat{\theta} + v$ is closer to the true value than $\hat{\theta} + w$ in the sense of first order stochastic dominance. Thus, values close to the estimated parameter value are more likely to be close to the true value than values that are further away. The CSbootstrap uses only the values that are close to the estimated parameter value, while the AD-bootstrap also uses the values that are further away. This gives an intuition as to why the CS-bootstrap yields confidence sets that are consistent in cases where the ADbootstrap does not yield such sets. Note that most of the discussion of confidence sets in the literature is about the coverage probability and about the length of the confidence set. This lemma and our examples add another consideration to the discussion of confidence sets in general. We now turn to investigating the differences between the AD-bootstrap and the CS-bootstrap within the context of two empirical studies.

4. A Comparison of the Confidence Sets Produced by the AD-bootstrap and the CS-bootstrap in Two Empirical Studies

In this section we use the parameter estimates and data from two empirical studies to compare the length of the confidence sets produced by the different methods discussed above. We first use results and data from Ham, Li and Shore-Sheppard (2011, hereafter HLSS). They estimate a model of the employment dynamics of disadvantaged mothers (i.e. single mothers with a high school degree or less) for the U.S. Specifically, they estimate hazard functions for these women for i) nonemployment spells in progress at the start of the sample, i.e. left censored nonemployment spells, ii) employment spells in progress at the start of the sample, i.e. left censored employment spells, iii) nonemployment spells that begin after the start of the sample, i.e. fresh nonemployment spells and iv) employment spells that begin after the start of the sample, i.e. fresh employment spells.²¹

 $^{^{21}}$ They also estimate the joint distribution of the (correlated) unobserved heterogeneity components in each hazard function.

HLSS first consider the effect of a change in an independent variable on the expected duration of each type of spell. Since the expected duration is a relatively simple differentiable function of the estimated parameters, they use the delta method to calculate confidence sets. In Table 1, we compare the confidence sets produced by the delta method, the AD-bootstrap, the CS-bootstrap and the WCS-bootstrap for these expected durations. Of these methods, only the AD-bootstrap may be inconsistent in the sense that no consistency proof exists. The first panel presents (for each type of spell) the confidence sets for the sample average of the individual expected durations for each spell type. The remaining panels show the analogous confidence sets (produced by each method for each type of spell) of the effect on the expected durations of i) having more schooling, ii) being African-American versus being white, iii) being Hispanic versus being white, and iv) having a child under 6 years versus not having a child under 6 years. For ease of viewing, in each panel we also report the ratio of the confidence interval lengths produced by: i) the AD-bootstrap relative to the delta method, ii) the CS-bootstrap relative to the delta method, and iii) the WCS-bootstrap relative to the delta method. From Table 1 we conclude that: i) the inconsistent AD-bootstrap produces somewhat shorter confidence sets than the delta method, ii) the CS-bootstrap produces substantially larger confidence sets than the delta method, and iii) the WCS-bootstrap produces, on average, confidence sets that are somewhat larger than those produced using the delta method but considerably smaller than those produced by the CS-bootstrap.

HLSS also consider the effect of the change in an independent variable on the estimated fraction of time a woman will spend in employment 3 years, 6 years, and 10 years after the change, which depends on the parameters from all the hazard functions. This function is nondifferentiable, so the delta method is no longer applicable and the CS-bootstrap or WCS-bootstrap should be used for calculating confidence sets.²² The first panel of

 $^{^{22}}$ For example, if a woman starts the sample in nonemployment, they calculate her hazard function for month 1 of a left censored nonemployment spell, and draw a uniform random number from [0,1]. Suppose the random number is less than the hazard. Then, she moves to employment and a 1 is registered for this month of her simulated employment history. In the next month, they calculate her hazard for month 1 of a fresh employment spell, and again draw a random number. If this random number is less than the hazard, a 0 is registered for the second month of her simulated employment history as she moves back to unemployment; otherwise, a 1 is registered for this month of her employment history as she stays in employment. This simulation is comparable to those used in structural modelling to estimate the effect of counterfactual policy changes.

Table 2 shows confidence sets for the baseline fraction of time spent in employment at 3 years, 6 years, and 10 years after the start of the sample. In the remaining panels, we show the respective confidence sets for the effects of changes in the demographic variables considered above on the fraction of time employed 3, 6, and 10 years after the change. In each case we also show the ratio of the confidence interval lengths produced by i) the AD-bootstrap relative to the CS-bootstrap and ii) the AD-bootstrap relative to the WCS-bootstrap. Table 2 shows that the CS and WCS-bootstrap confidence sets are basically identical, while the AD-bootstrap produces substantially smaller confidence sets than the consistent WCS-bootstrap and CS-bootstrap.

Finally, Lee and Ham (2012, here after LH) use data from an online dating service that proposes (opposite gender) matches to its individual members. The data indicate whether the man and woman agree to the date proposed by the company, and, if not, whether the man, the woman, or both turned down the date. The data set also contains information on whether, conditional on a first date, the couple goes on a second date, and, if not, whether the man, the woman, or both turned down the second date. Finally, the data also indicate whether the couple marries. Denote the outcome that individual i of gender j(j = M, F) accepts (refuses) date d (d = 1, 2) as $Y_d^j = 1 (Y_d^j = 0)$, and let the outcome where the couple marries (does not marry) be denoted by $Y^3=1$ ($Y^3=0$). LH estimate a fairly rich model of marriage and dating, and then simulate their estimated model to measure the relative efficiency of different possible matching algorithms that the dating company could use. Here we focus on the baseline probabilities of acceptance for the algorithm that the company actually uses. These probabilities are complicated differentiable functions of the estimated parameters, so it is sensible to use the CS-bootstrap to calculate confidence sets the baseline probabilities. In Table 3, we contrast these confidence sets with those produced by the AD-bootstrap. We find that the confidence sets produced by the ADbootstrap are about half the length of those produced by the CS-bootstrap, but that the CS-bootstrap still produces quite narrow confidence sets for the baseline probabilities.

Thus, our results suggest that previous work is likely to have substantially overstated the precision of their counterfactual policy effects, and that there may well be a significant efficiency gain from moving from the CS-bootstrap to the WCS-bootstrap.

5. Conclusion

Applied researchers often need to calculate confidence sets for functions of parameters that are nondifferentiable, or have unbounded or zero derivatives. Currently, they use the (nonparametric) bootstrap or sample from the asymptotic distribution of the estimators, since the delta method is not appropriate in these settings. Researchers also frequently use these procedures to obtain confidence sets for well-behaved, but complicated, functions. Indeed, two heavily cited articles and four prominent graduate econometrics textbooks recommend one or both of these approaches. Further, one of these approaches can be implemented using pre-programmed commands in the widely used Stata software package.

We first show that both of these procedures produce confidence sets that can be incorrect in the sense that the asymptotic coverage is less than intended, i.e. they produce confidence sets that are too small. We then propose two procedures that have correct coverage asymptotically under relatively weak conditions. In particular, our procedures are the first to give confidence sets for functions of parameters without restricting the derivatives of the functions and without requiring the functions to be continuous. We use data and parameter estimates from two empirical studies to compare our approach to the traditional one, and find that the procedures currently used produce substantially downward biased confidence sets.

Further, Andrews (2000) gives an example in which all versions of the bootstrap fail to consistently estimate the distribution of the maximum likelihood estimator. An example involving a function of a parameter yields the same maximum likelihood estimator as Andrews (2000). Our proposed procedures also work for this example, suggesting that it might be more fruitful to focus on the construction of confidence sets, rather than on the distributions of various versions of the bootstrap. Also, our methods yield confidence sets with asymptotically correct coverage for a class of models where the function of the parameters is not differentiable in the parameters of the data distribution, so that we complement the negative results of Hirano and Porter (2012).

Finally, one of our procedures (the WCS-bootstrap) produces asymptotically the same

confidence set as the delta method if the linear approximation holds, so in principle there is no efficiency loss in using the WCS-bootstrap in any application. Moreover, we find that in practice this procedure produces confidence sets similar to the delta method in a situation where the latter is likely to be used.

References

- Andrews, D.W.K. (1987): "Consistency in Nonlinear Econometric Models: A Generic Uniform Law of Large Numbers," *Econometrica*, 55: 1465–1472.
- [2] Andrews, D.W.K. (1991): "Heteroskedasticity and Autocorrelation Consistent Covariance Matrix Estimation," *Econometrica*, 59: 817–858.
- [3] Andrews, D.W.K. (2000): "Inconsistency of the Bootstrap when a Parameter is on the Boundary of the Parameter Space," *Econometrica*, 68: 399-405.
- [4] Andrews, D. W. K. and Guggenberger, P. (2010): "Asymptotic Size and a Problem with Subsampling and the *m* out of *n* Bootstrap," *Econometric Theory*, 26: 426-468.
- [5] Andrews, D. W. K. and X. Cheng (2012): "Estimation and Inference with Weak, Semi-strong, and Strong Identification," *Econometrica*, 80: 2153-2211.
- [6] Andrews, D. W. K. and X. Cheng (2013): "Maximum Likelihood Estimation and Uniform Inference with Sporadic Identification Failure," *Journal of Econometrics*, 173: 36-56.
- [7] Cameron, A. C. and P. K. Trivedi (2005): Microeconometrics, Methods and Applications, Cambridge University Press, Cambridge.
- [8] Cameron, A. C. and P. K. Trivedi (2010): Microeconometrics Using Stata (Revised Edition), Stata Press, College Station, Texas.
- [9] Chaudhuri, S., P. Goldberg, and P. Jia (2006): "Estimating the Effects of Global Patent Protection in Pharmaceuticals: A Case Study of Quinolones in India," American Economic Review, 96: 1477-1514.

- [10] Chernozhukov, V., H. Hong and E. Tamer (2007): "Parameter Set Inference in a Class of Econometric Models," *Econometrica*, 75: 1243–1284.
- [11] Eberwein, Z., C. J. Ham and R. LaLonde (1997): "The Impact of Being Offered and Receiving Classroom Training on the Employment Histories of Disadvantaged Women: Evidence From Experimental Data," *Review of Economic Studies*, 64: 655-682.
- [12] Eckstein, Z. and K. Wolpin (1999): "Why Youths Drop Out of High School: The Impact of Preferences, Opportunities and Abilities," *Econometrica*, 67: 1295–1339.
- [13] Eberwein, C., J. Ham, and R. LaLonde (2002): "Alternative Methods of Estimating Program Effects in Event History Models," *Labour Economics*, 9: 249-278.
- [14] Fitzenberger, B., A. Osikominu and M. Paul (2010): "The Heterogeneous Effects of Training Incidence and Duration on Labor Market Transitions," Mimeo, Albert– Ludwigs–University Freiburg.
- [15] Gaure, S., K. Røed, K. and L. Westlie (2010): "Job Search Institutions and Job Match Quality," Mimeo, The Ragnar Frisch Centre for Economic Research.
- [16] Greene, W. (2012): Econometric Analyses, 7th edition, Prentice Hall, New York, NY.
- [17] Ghysels, E. and A. Hall (2002): "Editors' Introduction to Twentieth Anniversary Commemorative Issue of the Journal of Business and Economic Statistics," *Journal* of Business and Economic Statistics, 20: 1-4.
- [18] Hall, P. (1995): The Bootstrap and Edgeworth Expansion, Springer Series, New York, NY.
- [19] Halton, J. H. (1964), Algorithm 247: "Radical-inverse quasi-random point sequence," Communications of the Association for Computing Machinery, 7: 701-702.
- [20] Ham, J., D. Mountain and L. Chan (1997): "Time-of-Use Prices and Electricity Demand: Allowing for Selection Bias in Experimental Data," *The Rand Journal of Economics*, 28: 113-141.

- [21] Ham, J., X. Li and L. Shore-Sheppard (2011): "Correcting for Seam Bias when Estimating Discrete Variable Models, with an Application to Analyzing the Employment Dynamics of Disadvantaged Women in the SIPP," Mimeo, University of Maryland.
- [22] Hamilton, J. D. (1994): Time Series Analysis, Princeton University Press, Princeton NJ.
- [23] Hansen, L. P. (1982): "Large Sample Properties of Generalized Method of Moments Estimators," *Econometrica*, 50: 1029-1054.
- [24] Hirano, K. and J. Porter (2012): "Impossibility Results for Nondifferentiable Functionals," *Econometrica*, 80: 1769–1790.
- [25] Hitsch, G., A. Hortacsu and D. Ariely (2010): "Matching and Sorting in Online Dating," American Economic Review, 100: 130-163.
- [26] Hoderlein, S. and S. Mihaleva (2008): "Increasing the Price Variation in a Repeated Cross Section," *Journal of Econometrics*, 147: 316-325.
- [27] Horowitz, J. L. (1998): Semiparametric Methods in Econometrics, Springer-Verlag, New York, NY.
- [28] Horowitz, J. L. (2001): "The Bootstrap" in *Handbook of Econometrics*, Vol. 5, ed. by J. J. Heckman and E. Leamer. Amsterdam: North-Holland.
- [29] Ichimura, H. (1993). "Semiparametric Least Squares (SLS) and Weighted SLS Estimation of Single Index Models," *Journal of Econometrics*, 58: 71–120.
- [30] Imbens, G. W. and C. F. Manski (2004): "Confidence Intervals for Partially Identified Parameters," *Econometrica*, 72: 1845-1857.
- [31] Inoue, A., and L. Kilian (2013): "Inference on Impulse Response Functions in Structural VAR Models," *Journal of Econometrics*, 177: 1–13.
- [32] Keane, M. and K. Wolpin (2000): "Eliminating Race Differences in School Attainment and Labor Market Success," *Journal of Labor Economics*, 18: 614-652.

- [33] Krinsky, I. and A. L. Robb (1986): "On Approximating the Statistical Properties of Elasticities," *The Review of Economics and Statistics*, 68: 715-719.
- [34] Lee, Sooyhung and J. Ham (2012). "Dating, Marriage, and Matching Online," Mimeo, University of Maryland.
- [35] Merlo, A. and K. Wolpin (2009): "The Transition from School to Jail: Youth Crime and High School Completion Among Black Males," Mimeo, University of Pennsylvania.
- [36] Newey, W. K., and D. McFadden (1994): "Large Sample Estimation and Hypothesis Testing," in the *Handbook of Econometrics*, Vol. 4, ed. by R. F. Engle and D. MacFadden. North-Holland, Amsterdam.
- [37] Newey, W. K. and K. D. West (1987): "A Simple, Positive Semi-definite, Heteroskedasticity and Autocorrelation Consistent Covariance Matrix," *Econometrica*, 55: 703–708.
- [38] Politis, D. N., and J. P.Romano (1994): "Large Sample Confidence Regions Based on Subsamples under Minimal Assumptions," Annals of Statistics, 22: 2031-2050.
- [39] Røed, K. and L. Westlie (2011): "Unemployment Insurance in Welfare States: Soft Constraints and Mild Sanctions," Forthcoming in the Journal of the European Economic Association.
- [40] Romano, J. P. and A. M. Shaikh (2010): "On the Uniform Asymptotic Validity of Subsampling and the Bootstrap," Mimeo, University of Chicago.
- [41] Ryan, S. (2012): "The Costs of Environmental Regulation in a Concentrated Industry," *Econometrica*, 80: 1019–1061.
- [42] Runkle, D.E. (1987): "Vector Autoregressions and Reality," Journal of Business and Economic Statistics, 5: 437-442.
- [43] Sobol, I. M. (1967): "Distribution of Points in a Cube and Approximate Evaluation of Integrals," U.S.S.R. Computational Mathematics and Mathematical Physics, 7: 86–112.

- [44] Stoye, J. (2009): "More on Confidence Intervals for Partially Identified Parameters," *Econometrica*, 77: 1299-1315.
- [45] Weisberg, S. (2005): Applied Linear Regression, 3rd edition, Wiley, New York, NY.
- [46] Wooldridge, J. M. (2010) Econometric Analysis of Cross Section and Panel Data, 2nd, MIT Press, Cambridge MA.

6. Appendix

Example 1:

Note that the true value of μ is zero. Consider

$$\begin{split} P\left(0 \in \left[\sqrt{|\bar{X}_N|} - \frac{1.96}{2\sqrt{|\bar{X}_N|}} \frac{1}{\sqrt{N}}, \sqrt{|\bar{X}_N|} + \frac{1.96}{2\sqrt{|\bar{X}_N|}} \frac{1}{\sqrt{N}}\right]\right) \\ &= P\left(\sqrt{|\bar{X}_N|} - \frac{1.96}{2\sqrt{|\bar{X}_N|}} \frac{1}{\sqrt{N}} \le 0 \le \sqrt{|\bar{X}_N|} + \frac{1.96}{2\sqrt{|\bar{X}_N|}} \frac{1}{\sqrt{N}}\right) \\ &= P\left(-\frac{1.96}{2\sqrt{|\bar{X}_N|}} \frac{1}{\sqrt{N}} \le -\sqrt{|\bar{X}_N|} \le \frac{1.96}{2\sqrt{|\bar{X}_N|}} \frac{1}{\sqrt{N}}\right) \\ &= P\left(-\frac{1.96}{2} \frac{1}{\sqrt{N}} \le -|\bar{X}_N| \le \frac{1.96}{2} \frac{1}{\sqrt{N}}\right) \\ &= P\left(-\frac{1.96}{2} \frac{1}{\sqrt{N}} \le \bar{X}_N \le \frac{1.96}{2} \frac{1}{\sqrt{N}}\right) \\ &= P\left(-\frac{1.96}{2} \frac{1}{\sqrt{N}} \le \bar{X}_N \le \frac{1.96}{2} \frac{1}{\sqrt{N}}\right) \\ &= \Phi\left(\frac{1.96}{2}\right) - \Phi\left(-\frac{1.96}{2}\right) \approx 0.67. \end{split}$$

Example 2

Consider $\rho = 1$. In that case $\hat{\beta} = \hat{\gamma}$, so that $h(\hat{\beta}, \hat{\gamma}) = h(\hat{\beta}, \hat{\beta})$. Therefore, we can define a new function that has just one scalar as its argument. In particular, define

$$\underline{h}(\beta) = \frac{1}{2}\Phi(\beta) + \frac{1}{2}\Phi\left(-2\beta - \sqrt{2\ln(2)}\right), \text{ and its derivative,}$$
$$\underline{h}'(\beta) = \frac{1}{2}\phi(\beta) - \phi\left(-2\beta - \sqrt{2\ln(2)}\right).$$

Note that

$$\underline{h}'(\beta = 0) = \frac{1}{2}\phi(0) - \phi\left(-\sqrt{2\ln(2)}\right)$$
$$= \frac{1}{2}\frac{1}{\sqrt{2\pi}} - \frac{1}{\sqrt{2\pi}}\exp\left[-\frac{1}{2}\left\{-\sqrt{2\ln(2)}\right\}^2\right]$$
$$= \frac{1}{2}\frac{1}{\sqrt{2\pi}} - \frac{1}{\sqrt{2\pi}}\exp\{-\ln(2)\} = 0.$$

Checking the second order conditions and the limits yields that $\underline{h}(0)$ is the minimum. Thus, $\underline{h}(\beta) > \underline{h}(0)$ for any $\beta \neq 0$. Therefore, the true value $\underline{h}(0) = \frac{1}{4} + \frac{1}{2}\Phi\left(-\sqrt{2\ln(2)}\right)$ is outside any two-sided AD-confidence interval of $\underline{h}(\beta)$. Thus, the coverage probability is zero in this case. Hence, the coverage probability is also zero for the function $h(\beta, \gamma) = \frac{1}{2}\Phi(\beta) + \frac{1}{2}\Phi\left(-2\gamma - \sqrt{2\ln(2)}\right)$ if $\rho = 1$. Note that the coverage probability is continuous in ρ so that the coverage probability is also too low for some $\rho < 1$. In the simulations, based on 100,000 repetitions, the coverage probability was still too low for $\rho = 0.5$.

Example 3:

Note that

$$P\left(\mu \in \left[\max\left(0, \bar{X}_N - \frac{1.96}{\sqrt{N}}\right), \max\left(0, \bar{X}_N + \frac{1.96}{\sqrt{N}}\right)\right]\right)$$

$$= P\left(\max\left(0, \bar{X}_N - \frac{1.96}{\sqrt{N}}\right) \le \mu \le \max\left(0, \bar{X}_N + \frac{1.96}{\sqrt{N}}\right)\right)$$

$$\ge P\left(\max\left(0, \bar{X}_N - \frac{1.96}{\sqrt{N}}\right) \le \mu \le \bar{X}_N + \frac{1.96}{\sqrt{N}}\right)$$

$$= 1 - P\left(\max\left(0, \bar{X}_N - \frac{1.96}{\sqrt{N}}\right) > \mu\right) - P\left(\bar{X}_N + \frac{1.96}{\sqrt{N}} < \mu\right)$$

$$= 1 - P\left(\max\left(-\bar{X}_N, -\frac{1.96}{\sqrt{N}}\right) > \mu - \bar{X}_N\right) - P\left(\frac{1.96}{\sqrt{N}} < \mu - \bar{X}_N\right).$$

Notice that $P\left(-\bar{X}_N > \mu - \bar{X}_N\right) = 0$ since $\mu \ge 0$. Thus

$$P\left(\mu \in \left[\max\left(0, \bar{X}_N - \frac{1.96}{\sqrt{N}}\right), \max\left(0, \bar{X}_N + \frac{1.96}{\sqrt{N}}\right)\right]\right)$$
$$\geq 1 - P\left(-\frac{1.96}{\sqrt{N}} > \mu - \bar{X}_N\right) - P\left(\frac{1.96}{\sqrt{N}} < \mu - \bar{X}_N\right)$$
$$= P\left(-1.96 \le \sqrt{N}(\bar{X}_N - \mu) \le 1.96\right)$$
$$= 0.95$$

since $\sqrt{N}(\bar{X}_N - \mu)$ has a standard normal distribution. This holds for any $\mu \ge 0$, including $\mu = \frac{c}{\sqrt{N}}$.

Proof of Theorem:

Consider a uniformly continuous function $f(\theta)$ and let f([0,1]) denote the set of values of $f(\theta)$ where $\theta \in [0,1]$, i.e.

$$f([0,1]) = \{ \tau \in \mathbb{R} | \tau = f(\theta) \text{ for some } \theta \in [0,1] \}.$$

Next, consider approximating this function on the interval $\theta \in [0,1]$ by evaluating the function at all the values of the M grid points, $G_M = \{\frac{1}{M}, ..., \frac{M-1}{M}, \frac{M}{M}\}$, and including all values that are no farther than $\eta > 0$ from $f(\frac{1}{M}), ..., f(\frac{M-1}{M})$, or $f(\frac{M}{M})$. The next lemma proves that this approximation contains the set f([0, 1]).

Lemma A1:

Let $f : \mathbb{R} \to \mathbb{R}$ be uniformly continuous. Let $\theta \in [0, 1]$, and $\eta > 0$. Then

$$f([0,1]) \subset \lim_{M \to \infty} \bigcup_{i=1}^{M} [f(i/M) - \eta, f(i/M) + \eta].$$

Proof: By construction, since η is fixed and $h(\theta)$ is uniformly continuous, there exists an r > 0 such that $y \in B_r(i/M)$ implies that $|f(y) - f(i/M)| < \eta$ where B_r denotes a ball with radius r. Thus, $f(B_r(i/M)) \subset [f(i/M) - \eta, [f(i/M) + \eta]$. Next, let $M > \frac{1}{r}$ so that $\bigcup_{i=1}^{M} B_r(i/M) = [0, 1]$. Finally, $f([0, 1]) \subset \bigcup_{i=1}^{M} f(B_r(i/M)) \subset \bigcup_{i=1}^{M} [f(i/M) - \eta, [f(i/M) + \eta]$.

Note that this lemma can easily be generalized to $\theta \in [0, 1]^2$ as well as $\theta \in [0, 1]^K$ or $\theta \in \Theta$, which is compact. Using this lemma, we now turn to the assumptions of the theorem. We first consider the case where Assumptions 1-2 hold and we use the CS-bootstrap. The vector-function $h(\theta)$ is uniformly continuous on Θ_r , r = 1, ..., R, so that for any $\eta > 0$ there is an $\varepsilon > 0$ such that $||h(\theta_1) - h(\theta_2)|| < \eta$ for all $\theta_1, \theta_2 \in \Theta_r$ with $||\theta_1 - \theta_2|| < \varepsilon$ where ||.||is the Euclidean norm. Therefore, we can partition the confidence set $CS_{1-\alpha}^{\theta}$ into Q sets, $CS_{1-\alpha}^{\theta}(1), CS_{1-\alpha}^{\theta}(2), ..., CS_{1-\alpha}^{\theta}(Q)$ such that (i) if $\theta_a \in CS_{1-\alpha}^{\theta}(q)$ and $\theta_b \in CS_{1-\alpha}^{\theta}(q)$ for some q, then $||h(\theta_a) - h(\theta_b)|| < \eta$; and (ii) $CS^{\theta}_{1-\alpha}(1) \cup CS^{\theta}_{1-\alpha}(2) \dots \cup CS^{\theta}_{1-\alpha}(Q) = CS^{\theta}_{1-\alpha}(Q)$ where $Q < \infty$. Note that such a partition is possible since Θ is compact. Also note that, without loss of generality, $CS^{\theta}_{1-\alpha}(1), CS^{\theta}_{1-\alpha}(2), ..., CS^{\theta}_{1-\alpha}(Q)$ have a nonzero Lebesgue measure. Thus, for any $M \ge M_0$, where $M_0 < \infty$, we have that every set $CS^{\theta}_{1-\alpha}(1)$, $CS_{1-\alpha}^{\theta}(2), ..., CS_{1-\alpha}^{\theta}(Q)$ has one or more of the grid point as its elements since the grid is dense in $CS^{\theta}_{1-\alpha}$. Thus, calculating $h(\theta_1), ..., h(\theta_M)$ and including every point in the image of $h(\theta), \theta \in \Theta$, that are no farther than $\eta > 0$ away from $h(\theta_1), ..., h(\theta_{M-1})$, or $h(\theta_M)$ gives $CS_{1-\alpha}^{h(\theta)} \subset CS_{1-\alpha}^{h(\theta)}$ for any $M \ge M_0$. Note that M_0 does not depend on N. Therefore, the requirement in Assumption 1,

$$\lim_{N \to \infty} \inf_{P \in \mathcal{P}} \Pr(\theta_0 \in CS^{\theta}_{1-\alpha}) \ge 1 - \alpha,$$

yields

$$\lim_{M \to \infty} \lim_{N \to \infty} \inf_{P \in \mathcal{P}} \Pr(h(\theta) \in \widehat{CS^{h(\theta)}_{1-\alpha}}) \ge 1 - \alpha.$$

Next, consider the case where Assumptions 2-3 hold and the researcher uses the WCSbootstrap and samples from the asymptotic distribution of $\hat{\theta}$. Note that by Assumption $3 w_k \neq 0, \ \hat{w}_k \neq 0, \ \sup_{P \in \mathcal{P}} |\hat{w}_k - w_k| = o_p(1)$. Also note that

$$WCS_{1-\alpha}^{\theta} = \{\theta \in \Theta | N \cdot (\hat{\theta} - \theta)' \hat{w} (\hat{w}' \hat{\Omega} \hat{w})^{-1} \hat{w}' (\hat{\theta} - \theta) \le \chi_{1-\alpha}^2 (H) \}.$$

Just as we could partition $CS_{1-\alpha}^{\theta}$, we can also partition $WCS_{1-\alpha}^{\theta}$ (since Θ is compact). Thus, we partition the confidence set $WCS_{1-\alpha}^{\theta}$ in Q sets,

 $WCS_{1-\alpha}^{\theta}(1), WCS_{1-\alpha}^{\theta}(2), ..., WCS_{1-\alpha}^{\theta}(Q)$ such that (i) if $\theta_a \in WCS_{1-\alpha}^{\theta}(q)$ and $\theta_b \in WCS_{1-\alpha}^{\theta}(q)$ for some q, then $||h(\theta_a) - h(\theta_b)|| < \eta$; and (ii) $WCS_{1-\alpha}^{\theta}(1) \cup WCS_{1-\alpha}^{\theta}(2) ... \cup WCS_{1-\alpha}^{\theta}(Q) = WCS_{1-\alpha}^{\theta}(\alpha)$, where $Q < \infty$. Also note that, without loss of generality, $WCS_{1-\alpha}^{\theta}(1), WCS_{1-\alpha}^{\theta}(2), ..., WCS_{1-\alpha}^{\theta}(Q)$ have a nonzero Lebesgue measure. Next, note that $\sqrt{N}(\hat{\theta} - \theta_0) \stackrel{d}{\rightarrow} N(0, \Omega)$ uniformly in $P \in \mathcal{P}$ so that a value of each of the Q subsets is sampled with probability approaching one as $M \to \infty$. Therefore, calculating $h(\tilde{\theta}_1), ..., h(\tilde{\theta}_M)$ and including every point in the image of $h(\theta), \theta \in \Theta$, that are no farther than $\eta > 0$ away from $h(\tilde{\theta}_1), ..., h(\tilde{\theta}_{M-1})$, or $h(\tilde{\theta}_M)$ gives $WCS_{1-\alpha}^{h(\theta)} \subset WCS_{1-\alpha}^{h(\theta)}$ with probability approaching one as $M \to \infty$. This yields the result for sampling from the asymptotic distribution.

Finally, consider the case where Assumptions 2-4 hold. In this case, one can use any version of the bootstrap as long as Assumption 4 is satisfied, i.e.

$$\lim_{N \to \infty} \inf_{P \in \mathcal{P}} \Pr(\hat{w}' \theta_0 \in WCS^{\theta}_{1-\alpha}) \ge 1 - \alpha.$$

Using the same reasoning as for sampling from the asymptotic distribution concludes the proof of the theorem.

Proof of Lemma:

Note that v is a constant. Thus, if θ , v, and w are scalars, then

$$P(|\hat{\theta} + v - \theta_0| \le \varepsilon) = P(|Z\sigma + v| \le \varepsilon),$$

where Z is a realization from a standard normal distribution. Note that this probability remains the same if v is replaced by (-v). Similarly, $P(|\hat{\theta} + w - \theta_0| \le \varepsilon)$ remains the same

if w is replaced by (-w). Thus, without loss of generality, we assume that $0 \le v < w$. This gives

$$P(|\hat{\theta} + v - \theta_0| \le \varepsilon) = P(-\varepsilon \le Z\sigma + v \le \varepsilon)$$
$$= P(-\frac{\varepsilon + v}{\sigma} \le Z \le \frac{\varepsilon - v}{\sigma}) = \int_{(-\varepsilon - v)/\sigma}^{(\varepsilon - v)/\sigma} \phi(z) dz.$$

Similarly,

$$P(|\hat{\theta} + w - \theta_0| \le \varepsilon) = \int_{(-\varepsilon - w)/\sigma}^{(\varepsilon - w)/\sigma} \phi(z) dz$$

This gives

$$\begin{split} P(|\hat{\theta} + v - \theta_0| \le \varepsilon) - P(|\hat{\theta} + w - \theta_0| \le \varepsilon) &= \int_{(-\varepsilon - v)/\sigma}^{(\varepsilon - v)/\sigma} \phi(z) dz - \int_{(-\varepsilon - w)/\sigma}^{(\varepsilon - w)/\sigma} \phi(z) dz \\ &= \int_{(\varepsilon - w)/\sigma}^{(\varepsilon - v)/\sigma} \phi(z) dz - \int_{(-\varepsilon - w)/\sigma}^{(-\varepsilon - v)/\sigma} \phi(z) dz \end{split}$$

using $0 \le v < w$. Note that the last equation holds, even if $\varepsilon - w < -\varepsilon - v$. Thus

$$\begin{split} P(|\hat{\theta} + v - \theta_0| \le \varepsilon) - P(|\hat{\theta} + w - \theta_0| \le \varepsilon) &= \int_{-w/\sigma}^{-v/\sigma} \phi(z + \varepsilon) dz - \int_{-w/\sigma}^{-v/\sigma} \phi(z - \varepsilon) dz \\ &= \int_{-w/\sigma}^{-v/\sigma} \{\phi(z + \varepsilon) - \phi(z - \varepsilon)\} dz. \end{split}$$

Note that $0 \le v < w$ so that $z \in [-w/\sigma, -v/\sigma)$ is negative. Also note that $\varepsilon > 0$ so that $\phi(z + \varepsilon) - \phi(z - \varepsilon) > 0$ for any $z \in [-w/\sigma, -v/\sigma]$. Therefore, $P(|\hat{\theta} + v - \theta_0| \le \varepsilon) - P(|\hat{\theta} + w - \theta_0| \le \varepsilon) > 0$. This completes the proof.

WCS-bootstrap and the delta method

Here we show that the WCS-bootstrap and the delta method are asymptotically equivalent under the standard assumptions of the delta method. The standard assumptions²³ of the delta method are (i) $\sqrt{N}(\hat{\theta} - \theta_0) \xrightarrow{d} N(0, \Omega)$, (ii) $\hat{\Omega} = \Omega + o_p(1)$, (iii) $h(\theta)$ is continuously differentiable in a neighborhood of θ_0 ; let $h_{Der}(\theta)$ denote this derivative and let $h_{Der} = h_{Der}(\theta_0)$. Let all elements of h_{Der} be nonzero. Let

$$CS_{1-\alpha}^{Delta} = \{h \in \mathbb{R}^H | h = h(\theta) \text{ for some } \theta \text{ for which }$$

$$N \cdot \{h(\hat{\theta}) - h\}' \{h_{Der}(\hat{\theta})' \hat{\Omega} h_{Der}(\hat{\theta})\}^{-1} \{h(\hat{\theta}) - h\} \le \chi^2_{1-\alpha}(H)\}.$$

 $^{^{23}}$ See, for example, Greene (2012, page 1084).

The coverage of this confidence set converges to $(1 - \alpha)$ under the assumptions stated above. We now show that the WCS-bootstrap yields the same confidence set asymptotically. Consider the confidence set for the WCS-bootstrap as $M \to \infty$,

$$\widehat{WCS_{1-\alpha}^{h(\theta)}} = \{h \in \mathbb{R}^H | h = h(\theta) \text{ for some } \theta \text{ for which } N \cdot (\hat{\theta} - \theta)' \hat{w} (\hat{w}' \hat{\Omega} \hat{w})^{-1} \hat{w}' (\hat{\theta} - \theta) \le \chi_{1-\alpha}^2 (H) \}.$$

First, consider the case that $h(\theta)$ is a linear function of the parameters so that $h(\hat{\theta}) - h(\theta) = w'(\hat{\theta} - \theta)$, and $\hat{w} = h_{Der}(\hat{\theta}) = w$. This gives

$$CS_{1-\alpha}^{Delta} = \{h \in \mathbb{R}^{H} | h = w'\theta \text{ for some } \theta \text{ for which } N \cdot (\hat{\theta} - \theta)' w (w'\hat{\Omega}w)^{-1} w'(\hat{\theta} - \theta) \le \chi_{1-\alpha}^{2}(H) \},$$
which is the same set as $\widehat{WCS_{1-\alpha}^{h(\theta)}}$.

Next, if $h(\theta)$ is continuously differentiable (and not necessarily linear), then $h(\hat{\theta}) - h(\theta) = h_{Der}(\bar{\theta})'(\hat{\theta} - \theta)$ where $\bar{\theta}$ is an intermediate value, $\bar{\theta} \in (\hat{\theta}, \theta)$. Note that $h_{Der}(\hat{\theta}), h_{Der}(\bar{\theta})$, and \hat{w} (calculated using numerical differentiation or least squares) all converge in probability to $w = h_{Der} = h_{Der}(\theta_0)$. This gives

$$CS_{1-\alpha}^{Delta} = \{h \in \mathbb{R}^H | h = w'\theta \text{ for some } \theta \text{ for which } N \cdot (\hat{\theta} - \theta)' w(w'\hat{\Omega}w)^{-1} w'(\hat{\theta} - \theta) \le \chi_{1-\alpha}^2(H) + o_p(1) \}$$

so that the confidence sets of the delta method and WCS-bootstrap are first order equivalent.

		1 2	1 5	1			
		Left-censored non-employment spells	Left-censored employment spells	Fresh non- employment spells	Fresh employment spells		
	Estimate	39.305	42.248	11.821	11.929		
Estimated Expected	Delta Method	[37.872,40.738]	[41.055,43.441]	[10.811,12.832]	[10.969,12.900]		
Estimated Expected Duration (in months)	AD-bootstrap	[38.009,40.491]	[40.957,43,327]	[10.897,12.884]	[11.031,12.965]		
	CI-bootstrap	[36.623,41.566]	[39.956,44.252]	[10.207,14.037]	[10.366,14.163]		
	WCI-bootstrap	[37.431,41.089]	[40.571,43.567]	[10.864,12.943]	[10.960,13.037]		
Ratio of Lengths of Confidence Intervals	AD/Delta	0.866	0.993	0.983	1.002		
	CI/Delta	1.725	1.801	0.528	1.966		
	WCI/Delta	1.276	1.256	0.972	1.076		
Effect on Expected Duration From Changes With Respect to:							
	T . 1 T 00			· · · · ·			

 Table 1: 95% Confidence Intervals for the Effects of Changes in the Demographic Variables (Separately) on the Expected

 Durations of Employment and Non-employment Spells

5.095 1.027 Estimated Effect 7.471 -0.070 [5.376,9.566] [3.142,7.048] [0.184, 1.870]Delta Method [-1.113,0.972] Age: [5.399,9.429] [3.209,7.057] [-1.094,0.919] [0.206, 1.825]AD-bootstrap (age=35) - (age=25) CI-bootstrap [3.550,10.945] [1.312,9.066] [-2.527, 2.066][-0.723, 2.655][0.024,2.037] WCI-bootstrap [5.167,9.623] [2.992,7.397] [-1.315,1.095] AD/Delta 0.962 0.985 0.965 0.960 **Ratio of Lengths of** 1.765 1.985 2.203 2.004 *CI/Delta* **Confidence Intervals** 1.128 1.156 1.194 WCI/Delta 1.063 **Estimated Effect** -5.293 7.013 -1.9402.970 [4.775,9.251] [-3.106,-0.773] [1.983,3.958] [-7.352, -3.234]Delta Method Schooling: AD-bootstrap [-7.230, -3.237] [4.795,9.211] [-3.120,-0.766] [2.013,3.940] (s = 12) - (s < 12)[-9.432,-1.204] CI-bootstrap [2.372,11.364] [-4.116,0.313] [1.099,5.011] WCI-bootstrap [-7.868, -2.615] [4.462,9.735] [-3.227, -0.600][1.823,4.057] AD/Delta 0.970 1.072 1.009 0.976 **Ratio of Lengths of** CI/Delta 1.998 2.184 1.898 1.981 **Confidence Intervals** WCI/Delta 1.276 1.280 1.126 1.131

Table 1 (Continued)								
		Left-censored	Left-censored	Fresh non-	Fresh			
		non-employment	employment	employment	employment			
		spells	spells	spells	spells			
Effect on Expected Duration From Changes With Respect to:								
	Estimated Effect	2.524	-1.074	1.842	-0.440			
Race:	Delta Method	[-0.022, 5.069]	[-3.390,1.242]	[0.434,3.249]	[-1.595,0.716]			
Black - White	AD-bootstrap	[0.100,5.265]	[-3.420,1.235]	[0.424,3.150]	[-1.550,0.695]			
	CI-bootstrap	[-1.702,7.678]	[-5.662,3.440]	[-0.752,4.612]	[-2.663,1.701]			
	WCI-bootstrap	[-0.355,5.674]	[-3.908,1.648]	[0.329,3.331]	[-1.775,1.005]			
Datio of Longths of	AD/Delta	1.015	0.910	0.968	0.971			
Confidence Intervals	CI/Delta	1.842	1.788	1.665	1.888			
Confidence filter vals	WCI/Delta	1.184	1.091	1.066	1.203			
	Estimated Effect	2.708	-2.616	0.435	0.090			
D	Delta Method	[-0.051,5.467]	[-5.623,0.391]	[-1.076,1.947]	[-1.330,1.511]			
Kace: Hispania Whita	AD-bootstrap	[-0.0422,5.728]	[-5.216,0.093]	[-0.968,1.889]	[-1.234,1.477]			
mspanie - winte	CI-bootstrap	[-2.921,9.055]	[-7.613,2.476]	[-2.225,3.490]	[-2.702,3.003]			
	WCI-bootstrap	[-0.522,6.455]	[-6.314,1.289]	[-1.105,2.103]	[-1.522,1.998]			
	AD/Delta	1.046	0.883	0.945	0.954			
Confidence Intervals	CI/Delta	2.170	1.678	1.891	1.887			
Confidence filter vals	WCI/Delta	1.264	1.264	1.061	1.164			
	Estimated Effect	4.225	-1.965	1.151	0.165			
Number of children	Delta Method	[2.450,6.000]	[-3.982,0.053]	[0.011,2.290]	[-0.721,1.051]			
less than 6 years old	AD-bootstrap	[2.335,6.042]	[-3.853,-0.080]	[0.012,2.295]	[-0.715,1.027]			
(one - zero)	CI-bootstrap	[0.558,7.912]	[-5.525,1.700]	[-1.205,3.250]	[-1.903,1.653]			
	WCI-bootstrap	[1.894,6.451]	[-4.359,0.653]	[-0.158,2.439]	[-0.875,1.164]			
	AD/Delta	1.044	0.935	1.002	0.983			
Katio of Lengths of Confidence Intervals	CI/Delta	2.072	1.722	1.955	2.007			
	WCI/Delta	1.284	1.242	1.140	1.151			

		3-year Period	6-year Period	10-year Period		
	Estimate	0.431	0.439	0.449		
Estimated Expected Fraction of	AD-bootstrap	[0.414,0.449]	[0.421,0.459]	[0.431,0.470]		
Time in Employment	CI-bootstrap	[0.396,0.469]	[0.401,0.480]	[0.409,0.491]		
	WCI-bootstrap	[0.396,0.469]	[0.401,0.480]	[0.409,0.489]		
Ratio of Lengths of Confidence	AD/CI	0.479	0.481	0.476		
Intervals	AD/WCI	0.479	0.481	0.488		

 Table 2: 95% Confidence Intervals For the Effect of Changing Demographic Variables on the Expected Fraction of Time Spent in Employment for Different Time Horizons

Change on the Expected Fraction of Time Spent in Employment With Respect to:

	Estimated Effect	0.09	0.097	0.100	
Schooling: $(s = 12) - (s < 12)$	AD-bootstrap	[0.072,0.107]	[0.078,0.115]	[0.081,0.119]	
Schooling: $(3 - 12) - (3 < 12)$	CI-bootstrap	[0.053,0.125]	[0.058,0.134]	[0.061,0.137]	
	WCI-bootstrap	[0.053,0.126]	[0.058,0.136]	[0.061,0.140]	
Ratio of Lengths of Confidence	AD/CI	0.486	0.487	0.500	
Intervals	AD/WCI	0.479	0.474	0.481	
	Estimated Effect	-0.031	-0.034	-0.037	
Page: Plack White	Estimated Effect AD-bootstrap	-0.031 [-0.050,-0.009]	-0.034 [-0.055,-0.011]	-0.037 [-0.058,-0.012]	
Race: Black - White	Estimated Effect AD-bootstrap CI-bootstrap	-0.031 [-0.050,-0.009] [-0.073,0.019]	-0.034 [-0.055,-0.011] [-0.080,0.019]	-0.037 [-0.058,-0.012] [-0.083,0.019]	
Race: Black - White	Estimated Effect AD-bootstrap CI-bootstrap WCI-bootstrap	-0.031 [-0.050,-0.009] [-0.073,0.019] [-0.073,0.019]	-0.034 [-0.055,-0.011] [-0.080,0.019] [-0.079,0.019]	-0.037 [-0.058,-0.012] [-0.083,0.019] [-0.083,0.019]	
Race: Black - White Ratio of Lengths of Confidence	Estimated Effect AD-bootstrap CI-bootstrap WCI-bootstrap AD/CI	-0.031 [-0.050,-0.009] [-0.073,0.019] [-0.073,0.019] 0.446	-0.034 [-0.055,-0.011] [-0.080,0.019] [-0.079,0.019] 0.444	-0.037 [-0.058,-0.012] [-0.083,0.019] [-0.083,0.019] 0.451	

Table 2 (Continued)							
3-year Period 6-year Period 10-year Perio							
Change on the Expected Fraction of Time Spent in Employment With Respect to:							
	Estimated Effect	-0.026	-0.028	-0.029			
Daga: Hispania Whita	AD-bootstrap	[-0.046,-0.001]	[-0.050,-0.002]	[-0.052,-0.001]			
Kace. Inspanie - winte	CI-bootstrap	[-0.068,0.023]	[-0.073,0.024]	[-0.077,0.026]			
	WCI-bootstrap	[-0.070,0.025]	[-0.074,0.027]	[-0.077,0.029]			
Ratio of Lengths of Confidence	AD/CI	0.495	0.495	0.495			
Intervals	AD/WCI 0.474 0.4		0.475	0.481			
	Estimated Effect	-0.030	-0.033	-0.035			
Number of kids less than 6	AD-bootstrap	[-0.047,-0.014]	[-0.052,-0.016]	[-0.054,-0.017]			
years old: one - zero	CI-bootstrap	[-0.067,0.004]	[-0.072,0.003]	[-0.075,0.003]			
	WCI-bootstrap	[-0.067,0.004]	[-0.072,0.003]	[-0.075,0.003]			
Ratio of Lengths of Confidence	AD/CI	0.465	0.480	0.474			
Intervals	AD/WCI	0.465	0.480	0.474			

						Prediction	CI-bootstrap	AD-bootstrap	(2)length /(3)length
						(1)	(2)	(3)	(4)
Panel A									
Outcomes	Y_1^M	Y_1^W	Y_2^M	Y_2^W	Y_2				
1	0	0				57.425	[56.576,57.841]	[56.793,57.579]	1.609
2	0	1				10.758	[10.422,11.210]	[10.545,10.966]	1.872
3	1	0				16.111	[15.481,16.869]	[15.878,16.488]	2.275
4	1	1	0	0		4.354	[4.198,4.672]	[4.301,4.494]	2.456
5	1	1	0	1		2.393	[2.304,2.539]	[2.329,2.450]	1.942
6	1	1	1	0		3.181	[3.083,3.396]	[3.133,3.300]	1.874
7	1	1	1	1	0	5.524	[5.144,5.970]	[5.295,5.809]	1.607
8	1	1	1	1	1	0.255	[0.231,0.300]	[0.238,0.289]	1.353

Table 3: Estimated Probabilities of Different Dating Outcomes under the Company's Matching Algorithm